



ELBOW RECONSTRUCTIVE SURGERY PROTOCOL

Triceps Tendon Repair & Reconstruction

Clinical Rationale & Pathophysiology: Anatomical re-anchoring of ruptured distal triceps tendon to the olecranon process utilizing transosseous bone tunnels or double-row knotless suture bridge anchors. Protects healing enthesis from tensile failure during extension.

CRITICAL PRECAUTIONS & IMMOBILIZATION RULES

- Hinged elbow brace required at all times for 6 weeks (locked at 30°-45° initially).
- Brace unlock settings: Wks 1-2 locked 30°-45°; Wks 3-4 unlocked 0°-60°; Wks 5-6 unlocked 0°-90°; Wk 7+ full motion.
- STRICTLY NO ACTIVE RESISTED ELBOW EXTENSION or pushing up from chairs for 8 to 12 weeks.
- No lifting, carrying, or pushing with operative arm (>1 lb) for 12 weeks.

Phase I: Immobilization & Controlled Passive Motion (Weeks 0 - 6)

Rehabilitation Goals & Milestones	Therapeutic Interventions & Exercise Progression
<ul style="list-style-type: none">• Protect transosseous / anchor fixation at olecranon• Gradual protected passive flexion progression via weekly brace adjustments• Prevent elbow flexion/extension joint contracture	<ul style="list-style-type: none">• Hinged brace worn 24/7 except supervised exercises• Passive elbow flexion within safe envelope; extension blocked by brace• Active forearm pronation/supination and wrist/hand range of motion• Active shoulder, wrist, and hand exercises to maintain kinetic chain• Cryotherapy for post-surgical posterior elbow edema control

Phase II: Discontinue Brace & Active Extension (Weeks 6 - 12)

Rehabilitation Goals & Milestones	Therapeutic Interventions & Exercise Progression
<ul style="list-style-type: none">• Discontinue hinged brace at 6 weeks• Attain full unassisted active elbow flexion and active extension without resistance• Eliminate elbow extension lag without forceful joint manipulation	<ul style="list-style-type: none">• Active unassisted elbow flexion and gravity-assisted active extension• Full available active pronation, supination, and functional motion• Gentle passive terminal flexion and extension stretching (low load)• Light periscapular and shoulder strengthening• Initiate submaximal isometric triceps contraction at 60° at week 8

Phase III: Progressive Strengthening & Resistance (Weeks 12 - 16)

Rehabilitation Goals & Milestones	Therapeutic Interventions & Exercise Progression
<ul style="list-style-type: none">• Restore triceps strength and muscular endurance• Progress functional lifting to 10-15 lbs by week 16• Eliminate elbow extension lag	<ul style="list-style-type: none">• Begin light resisted triceps extension (1-2 lb dumbbell) and elastic bands• Progressive closed chain wall pushups and upper body ergometer• Overhead reaching and light functional triceps endurance drills• Progressive serratus anterior and periscapular strengthening

Phase IV: Unrestricted Return to Heavy Loading & Sport (Weeks 16 - 24)

Rehabilitation Goals & Milestones	Therapeutic Interventions & Exercise Progression
<ul style="list-style-type: none">• Full return to heavy manual labor, contact sports, and bodybuilding/lifting• Objective strength testing: elbow extension strength symmetry >85-90%	<ul style="list-style-type: none">• Sport-specific power drills, high-resistance triceps loading• Heavy bench press, dips, and overhead pressing after 5-6 months• Final surgeon clearance for unrestricted vocational/athletic duties

Clinical Communications & Triage: Division of Orthopaedic Surgery, UHN Toronto Western Hospital · Inquiries: office@drcvortho.ca · Protocol Portal: